

NOTE XXXIII.

ZOOLOGICAL RESEARCHES IN LIBERIA.

A LIST OF THE BRENTHIDAE,

collected by J. BÜTTIKOFER, C. F. SALA and
F. X. STAMPFLI

BY

Dr. ANGELO SENNA,

Assistant in the R. Museum at Florence.

The small collection of Brenthidæ obtained in Liberia by the well-known and successful travellers Büttikofer, Sala and Stampfli, and submitted to me by Mr. C. Ritsema, amounts to 27 specimens, all belonging to described species. Nevertheless I thought it useful to publish the following list as it throws some further light upon the distribution of the African Brenthidæ.

1. *Zemioses porcatus* Pascoe, Journ. of Entom. I, p. 394, 1862.

Six specimens collected by Büttikofer.

The locality of the type of this species is Port Natal, but recently Mr. L. Conradt obtained it in Togoland.

2. *Symmorphocerus semipunctatus* Pascoe, Ann. and Mag. of Nat. Hist. ser. 4, X, p. 321, pl. XV, figs. 7 and 7a, ♂, 1872 (sub *Cordus*).

One female collected by Büttikofer.

This species, originally described from Port Natal, is

Notes from the Leyden Museum, Vol. XVII.

now for the first time recorded from West Africa. It is closely allied to *S. Alluaudi* Senna ¹⁾ from Yoruba (Oyo), but distinguished by the apophyses of the metarostrum which have the sides oblique, not parallel. *Symmorphocerus frontalis* Oliv. ²⁾ from Guinea and Natal, which has the sides of the apophyses likewise oblique, differs by its larger size, by the different colour, and by the less punctured elytral striae.

3. *Spatherhinus gabonicus* Thomson, Arch. entom. II, p. 116, 1858 (sub *Arrhenodes*).

One male and one female from the Junk River, collected by Mr. Stampfli.

A common species, recorded from Assinie, Gaboon and Congo.

4. *S. longiceps* Kolbe, Entom. Nachrichte, XIV, p. 308, 1888.

One male collected by Büttikofer and Sala.

The habitat of this remarkable species is the same as that of the preceding one. Mr. Power had named this species *S. liberiae* Pow. in litt.

5.? *S. picturatus* Kolbe, Entom. Nachrichte, XIV, p. 306, 1888.

One female, in bad condition, from the Junk River, collected by Mr. Stampfli.

This species has been described from a single male from Cuango; Mr. Alluaud captured it in Assinie.

6. *Eupsalis gentilis* Thomson, Arch. entom. II, p. 117, 1858 (sub *Arrhenodes*).

Two males collected by Büttikofer and Sala.

A common West African species.

7. *Ceocephalus picipes* Olivier, Entom. V, n°. 84, p. 442, pl. 2, fig. 18, ♀, 1807—8.

1) Ann. Soc. Entom. France, LXIII, p. 409, fig. 1, 1894.

2) Entomologie, V, n°. 84, p. 434, pl. 2, fig. 14, 1807—8.

Several specimens from the Junk River and Cape Mount, collected by Büttikofer, Sala and Stampfli.

The commonest species of Ethiopical Brenthidæ.

8. *Storcosomus Rissi* Labram & Imhoff, Gen. Curc. I, n^o. 12, 1842.

One female from the Junk River, obtained by Mr. Stampfli.

Florence, March 11th 1896.